



Hertfordshire and
West Essex Integrated
Care System

PCN Insights Pack 2024

Epping Forest North

Stefania Mistretta, PHM Champion
Hannes van der Merwe, PHM Champion
Jaron Inward, Snr PHM Champion



Working together
for a healthier future

Introduction

This latest version of the PCN pack has been developed around the ICB Clinical Priorities signed off by the ICB Board in March 23 and how they align to the population outcomes we are aiming to achieve. (See tables on the right-hand side.)

The data contained within this pack compares the PCN data against Place and ICB. An overview table of all the data is available on the following page with areas of opportunity highlighted.

Some data will not be available at PCN level e.g. mortality rate for CVD. Where this is the case, proxy measures that will lead to improving this will be included e.g. early identification.

Where opportunities link with areas within the ECF or QOF a selection of related indicators have been shared alongside a link to Ardens Manager.

The data within this pack are shown at PCN level and are health focused. However, to improve outcomes for our population, input from many partners is required. This pack can facilitate discussion within your Integrated Neighbourhood Teams as described within the Primary Care Strategy and the wider HCP transformation spaces supported by your Primary Care Transformation leads.

Packs for other PCNs, Localities and Place can be found:

[Population health management – Hertfordshire and West Essex Integrated Care System \(hertsandwestessexics.org.uk\)](https://hertsandwestessexics.org.uk)

Here you can also find previous PCN packs outlining the descriptive demographics for the PCNs.



Area	Clinical Priority
CYP	<ul style="list-style-type: none"> Improved Readiness for school in children eligible for FSM Reduce rates of Childhood obesity Reduced unnecessary A&E attendances and admissions
Prevention and Health Inequalities	<ul style="list-style-type: none"> Reduced premature mortality rate for CVD
LTC & Frailty	<ul style="list-style-type: none"> Reduce attendance and admissions for falls, people with frailty and people in last year of life Development of more proactive, preventative care models for management of LTC and Frailty
Mental Health	<ul style="list-style-type: none"> Reducing suicide rates and attendances/ admission rates for self-harm Reducing rates of A&E attendances involving substance misuse and violence

Outcome	Definition
Improve life expectancy	Average age at death for people who have died in the last 12 months
Improve healthy life expectancy	Average age of people who have left the 'healthy' segment in the last 12 months
Reduce the proportion of people living with advanced disease and complexity	Proportion of the registered population who are in the advanced disease and complexity segment (segment)
Reduce the rate of ambulatory care sensitive emergency hospital admissions	Rate of ambulatory care sensitive emergency admissions in the last 12 months
Reduce the overall spend on emergency hospital admissions	Spend on Emergency hospital admissions within a financial year

Epping Forest North at a Glance

The table on the right provides a summary of the data contained within this update highlighting showing how the PCN data compares with Place and ICB.

For Epping Forest North areas highlighted are

- **Childhood Obesity at Year 6**
- **Observed versus expected prevalence of LTC**
- **% of people for secondary prevention CVD who are on low and medium intensity statins**
- **Admissions for Hip Fractures (75+)**
- **Prevalence of Learning Disabilities**

The following link takes you to Ardens Manager where there are reports. Here you will find the latest information on identification of LTCs and details of case finding Ardens searches available within EMIS and System one.

Clinical Priority	Metric	PCN compared to Place average	PCN compared to ICB average
Childhood obesity	% of children in Reception who are overweight	↓	↓
	% of children in Year 6 who are overweight	↑	↑
Reduce rates of emergency care for children and young people	A&E Attendances for Asthma (Children)	↓	↑
	Admissions for Asthma (Children)	↑	↔
	Admissions for Wheeze (Children)	↑	↓
	Admissions for Diabetes (Children)	↔	↔
	Admissions for Epilepsy (Children)	↔	↔
Prevention and health inequalities (Premature mortality for CVD)	Lifestyle risk factors: Smoking	↔	↔
	Observed versus expected prevalence	↓	↓
	Annual Reviews completed for LTCs	↑	↑
	% of people with AF treated with Anti Coagulant	↔	↔
	Control of hypertension	↔	↔
Preventative, Proactive care models for LTC	Identification of hypertension	↑	↑
	% of people for secondary prevention CVD who are on low and medium intensity statins	↑	↑
	% of people living with diabetes with all 8 care processes completed	↔	↓
	Reduction in prevalence of ACS conditions	↓	↓
Preventative, Proactive care models for frailty and EOL	Admissions for falls (75+)	↑	↓
	Admissions for Hip Fractures (75+)	↑	↑
Mental Health	Prevalence of Mental Health Conditions including LD	↓ (LD)	↓ (LD)
	Admissions for Self-Harm	↓	↓



<https://app.ardensmanager.com/login>

Rates of Childhood Obesity

CYP outcome – Every child will have the best start and live a healthy life

ICB overarching outcome of Improving Healthy life expectancy

- In keeping with the national data, the PCN rates for Childhood Obesity are higher for year 6 in comparison to reception children.
- Compared to the ICB and Place rate, Epping Forest North PCN has lower rates of Childhood Obesity for Children in Reception. However, the rate of Childhood Obesity at Year 6 is higher than both Place and ICB.
- The data suggest that there is a deterioration from reception to Year 6 in childhood obesity when comparing against Place and ICB.





A&E attendances for Asthma (CYP)

CYP outcome – Reduce the number of unplanned admissions for long term conditions

ICB overarching outcome of reduce the proportion of people living with advanced disease and complexity

- The data shows that Epping Forest North PCN has a higher rate of A&E attendances for Children and Young People with Asthma (rolling years data on the right-hand side) compared with the ICB rate, but lower than Place
- Rates of Children and Young People attending A&E for Asthma have increased since 2020/21 post covid, with the latest data similar to pre covid levels.
- The data should continue to be monitored to understand the trends.




Admissions for Asthma (CYP)

- Compared to Place, Epping Forest North PCN has a higher rate of children's admissions for Asthma (rolling years data on the right-hand side).
- There is variation in admissions between the practices with no admissions in the 12 months up to November 23 for Abridge.
- Higher Proportion of Asthma Reviews are carried out within Epping Forest North PCN in comparison with Place and the ICB, however the QOF is for all ages and children specific reviews cannot be identified within the data.
- The Children and Young Peoples programme can be contacted via hweicbenh.cypteam@nhs.net for details of current projects.


CYP outcome – Reduce the number of unplanned admissions for long term conditions

ICB overarching outcome of reduce the proportion of people living with advanced disease and complexity


Asthma Inpatient Activity DSR per 100,000 by GP for PCN: Epping Forest North and period Dec-22 to Nov-23 against PCN, Place & ICB rates



Asthma Inpatient Activity DSR per 100,000 by Year for PCN: Epping Forest North against ICB & Place rates



Percentage of patients with asthma, on the register, who have had an asthma review in the preceding 12 months: Epping Forest North PCN





Admissions for Wheeze (CYP)

CYP outcome – Reduce the number of unplanned admissions for long term conditions

ICB overarching outcome of reduce the proportion of people living with advanced disease and complexity

- Epping Forest North PCN has a higher rates of Children and Young People admitted to hospital for abnormalities of breathing including wheeze compared to Place but lower than place.
- Rates of Children and Young People admitted to Hospital for Wheeze fluctuate annually with the latest forecast outturn from November data showing a decrease on the previous year.
- When looking at the data by practice there is variation.





Admissions for Diabetes (CYP)

CYP outcome – Reduce the number of unplanned admissions for long term conditions

ICB overarching outcome of reduce the proportion of people living with advanced disease and complexity

- Diabetes is identified as an area of focus within the Core 20 plus 5 for children. The rate of admission for the PCN is similar to Place and ICB (latest 12 months data on the right-hand side).
- The numbers of children admitted for diabetes are small and this should be considered when looking at the data.
- The data for diabetes will continue to be monitored at wider HCP and ICB footprints.





Admissions for Epilepsy (CYP)

CYP outcome – Reduce the number of unplanned admissions for long term conditions

ICB overarching outcome of reduce the proportion of people living with advanced disease and complexity

- Epilepsy is identified as an area of focus within the Core 20 plus 5 for children. The rate of admission for the PCN is similar to Place and ICB (latest 12 months data on the right-hand side).
- The numbers of children admitted for diabetes are small and this should be considered when looking at the data.
- The data for diabetes will continue to be monitored at wider HCP and ICB footprints.




Prevention and health inequalities – Lifestyle factors - Smoking


- Epping Forest North PCN data for recording of smoking status shows a similar picture to the Place and ICB.
- A slightly higher proportion of patients have been offered treatment for smoking compared to place and ICB.
- The table to the right gives detail by condition of the opportunity for further recording of smoking status. This shows the position in January. The latest position can be found on <https://app.ardensmanager.com/login>

Practices	ECF 2023-24 - Condition Section Under Smoker, Smoking Status, and Smoking Status Recorded - as of Jan. 2024					
	Pre-Diabetes		Diabetes		Atrial Fibrillation	
	Remaining % of Population with a Smoking status	Smoking Available Patients - Number	Remaining % of Population with a Smoking status	Smoking Available Patients - Number	Remaining % of Population with a Smoking status	Smoking Available Patients - Number
Abridge Surgery	53%	223	29%	234	0%	6
High Street Surgery, Epping	28%	404	25%	452	0%	9
Maynard Court Surgery	40%	368	17%	703	0%	17
Ongar Health Centre	28%	1831	6%	822	0%	19
The Limes Medical Centre	33%	1040	24%	1259	0%	21
The Market Square Surgery	23%	660	9%	1109	0%	19


Percentage of patients* whose notes record smoking status in the preceding 12 months



Percentage of patients aged 15+ who are current smokers with a record of an offer of support & treatment within preceding 24 months



Percentage of patients* who are current smokers with a record of an offer of support & treatment within the preceding 12 months



Hertfordshire and West Essex Integrated Care System



* with any or any combination of the following conditions: CHD, PAD, stroke or TIA, hypertension, diabetes, COPD, CKD, asthma, schizophrenia, bipolar affective disorder or other psychoses

Prevention and health inequalities

Early Identification: Expected vs observed prevalence


LTC Outcome – Proportion of people with a long-term condition who feel able to manage their condition

ICB overarching outcome of reduce the proportion of people living with advanced disease and complexity

The data on this page shows the national modelled estimated prevalence for the PCN compared with the latest published QOF prevalence for the PCN.

- Epping Forest North PCN recorded prevalence compared with the modelled estimated prevalence for the PCN is lower across all conditions.
- The data shows an opportunity for further identification for these conditions. Case finding searches can be found within the Ardens Suite of searches. <https://app.ardensmanager.com/login>

Disease/ Condition	Disease Detection Modelling for Epping Forest North PCN - No. of New Diagnoses to Meet ICS & PLACE Rates - 2023/24	
	Number to meet ICS rate	Number to meet PLACE rate
Asthma	3832	70
Atrial Fibrillation		52
Chronic Kidney Disease	1802	217
Chronic Obstructive Pulmonary Disease		5
Coronary Heart Disease	1782	8
Diabetes Mellitus		46
Epilepsy	376	33
Heart Failure		1
Hypertension	9076	83
Non-Diabetic Hyperglycaemia		384
Peripheral Arterial Disease	286	12
Stroke and Transient Ischaemic Attack	1081	9



Development of more proactive, preventative care models for LTC - Prevalence

LTC Outcome – Reduce the proportion of people with a long-term condition who are in the advanced disease & complexity or frailty & end of life segments

ICB overarching outcome of reduce the proportion of people living with advanced disease and complexity

Following the expected and observed prevalence modelling in the previous slide, where there are opportunities, this slide looks at the recorded prevalence by practice for the PCN compared with Place and ICB.

The development of an ICB Data Platform will create a longitudinal record for our patients which will allow the data to be viewed by different characteristics such as deprivation, ethnicity, co-morbidities.

For LTCs, QOF and ECF data can help us understand areas of opportunity for improving outcomes for our population living with LTCs.

Epping Forest North PCN - Long-Term Conditions 2022-23 QOF Prevalence, with 3 Year Trend.

QOF Disease/ Condition	QOF 22-23 - ICB %	QOF 22-23 - PLACE %	QOF 22-23 - PCN %	THE LIMES MEDICAL CENTRE		ONGAR HEALTH CENTRE		HIGH STREET SURGERY, EPPING		ABRIDGE SURGERY		MAYNARD COURT SURGERY		MARKET SQUARE SURGERY	
				QOF 2022-23	3 Year Trend	QOF 2022-23	3 Year Trend	QOF 2022-23	3 Year Trend	QOF 2022-23	3 Year Trend	QOF 2022-23	3 Year Trend	QOF 2022-23	3 Year Trend
				Asthma	6.16%	6.29%	6.18%	6.11%	↗	6.37%	↗	5.90%	↘	6.06%	↘
Atrial fibrillation	2.09%	2.20%	2.42%	2.66%	↗	2.98%	↗	2.18%	↗	2.86%	↗	1.90%	↗	1.96%	↗
Chronic kidney disease	3.46%	3.61%	4.32%	2.65%	↗	6.24%	↘	7.06%	↗	1.35%	↗	4.16%	↘	4.47%	↗
Chronic obstructive pulmonary disease (COPD)	1.49%	1.54%	1.78%	1.67%	↗	1.69%	↘	1.53%	↘	1.54%	↘	2.54%	↗	1.69%	↘
Diabetes mellitus	6.63%	6.86%	7.33%	7.31%	↗	7.34%	↗	6.76%	↘	6.00%	↘	8.04%	↗	8.53%	↗
Epilepsy	0.70%	0.67%	0.61%	0.74%	↗	0.59%	↗	0.54%	↘	0.54%	↗	0.59%	↗	0.63%	↗
Heart Failure	0.80%	1.00%	1.42%	1.03%	↗	3.06%	↘	0.99%	↘	1.07%	↗	1.17%	↗	1.17%	↘
Hypertension	13.84%	14.64%	16.09%	14.40%	↗	18.82%	↗	13.77%	↗	17.78%	↗	15.80%	↗	15.99%	↗
Non-diabetic hyperglycaemia	6.42%	6.49%	7.92%	6.33%	↗	18.99%	↗	6.70%	↗	5.23%	↗	4.58%	↗	5.68%	↗
Peripheral arterial disease	0.44%	0.46%	0.46%	0.43%	↘	0.47%	↘	0.41%	↗	0.39%	↘	0.56%	↗	0.48%	↗
Secondary prevention of coronary heart disease	2.67%	2.80%	3.12%	2.80%	↗	3.80%	↘	2.67%	↘	3.37%	↘	3.10%	↗	2.97%	↘
Stroke and transient ischaemic attack	1.63%	1.64%	1.90%	2.06%	↗	2.49%	↗	1.57%	↘	1.93%	↘	1.62%	↘	1.73%	↘



Development of more proactive, preventative care models for LTC : Annual Reviews

- The table on the right shows a summary of the percentage of patients receiving an annual review or risk assessment by condition.
- Where the cell is highlighted, the percentage is lower than the place value.
- The data shows that the PCN has higher percentages of reviews completed compared to place and ICB apart from NDH reviews.
- The source of data in this table is QOF national reporting. More detailed information with the latest position is available to practices via <https://app.ardensmanager.com/login>

	ICB	WE	Epping Forest North PCN	Abridge Surgery	High Street Surgery, Epping	Market Square Surgery	Maynard Court Surgery	Ongar Health Centre	The Limes Medical Centre
% of AF Patients with Stroke Risk Assessed in the last 12 months	92.94	93.03	94.7	95.9	96.0	96.1	91.8	96.7	92.8
The % of patients aged 45 or over who have a record of blood pressure in the preceding 5 years	85.65	86.44	88.3	90.3	89.3	90.3	89.7	88.9	85.2
The % of patients with a diagnosis of heart failure on the register, who have had a review in the preceding 12 months	72.69	70.02	71.0	88.6	75.0	88.0	80.2	84.9	18.5
The % of patients with asthma, on the register, who have had an asthma review in the preceding 12 months	64.02	61.44	62.6	67.7	70.5	66.0	68.1	63.6	52.3
The % of patients with COPD, on the register, who have had a review in the preceding 12 months	75.74	74.01	76.9	84.1	83.8	88.4	83.0	76.1	60.4
The % of patients with diabetes, on the register, with a record of a foot examination and risk classification: 1) low risk (normal sensation, palpable pulses), 2) increased risk (neuropathy or absent pulses), 3) high risk (neuropathy or absent pulses plus deformity or skin changes in previous ulcer) or 4) ulcerated foot within the preceding 12 months	75.54	72.97	82.7	75.9	81.6	89.1	88.2	95.7	67.5
The % of patients with non-diabetic hyperglycaemia who have had an HbA1c or fasting blood glucose performed in the preceding 12 months.	82.92	80.46	76.6	82.2	90.2	90.0	86.6	69.6	71.6





Hertfordshire and West Essex Integrated Care System





Prevention and health inequalities – Atrial Fibrillation

- Epping Forest North PCN recorded prevalence for Atrial Fibrillation is higher than both Place and the ICB prevalence.
- Once identified the percentage of patients currently treated with anti-coagulant drug therapy is similar to the Place and ICB, however there is variation between the practices.
- The latest AF indicators can be found at <https://app.ardensmanager.com/login>



Prevention and health inequalities – Hypertension


- Epping Forest North PCN recorded prevalence for hypertension is higher than both Place and the ICB prevalence.
- Once identified the data shows the percentage of patients in whom the latest BP reading is below the age-appropriate treatment threshold is slightly higher than Place and ICB, however there is variation between the practices.
- The latest hypertension indicators can be found at <https://app.ardensmanager.com/login>




Lipid management: Percentage of people on lipid lowering therapy for secondary prevention who are on low and medium intensity statins

- National lipid management pathways ([Link to guidance](#)) recommend the use of high intensity statins for all people with a history of Cardiovascular disease as well as where high dose statins are needed to control cholesterol. People on high intensity statins will see a greater reduction in c-LDL levels and reduce the risk of cardiovascular events.
- Data from OpenPrescribing provides information on the proportion of people who are on statins that are currently prescribed low or medium intensity doses. The ICS is a negative outlier in this area, with a high proportion of people not on a high intensity statin.
- The data for the PCN shows that there is an opportunity to improve statin treatment, prescribing a higher proportion of people onto high intensity statins. The PCN is in the 96th percentile with 37.4% of people not on high intensity statins. This compares to 28.3% nationally.

PCN – Items of Low and Medium Intensity Statins as a Percentage of Items of All Statins by Year





ICB – Items of Low and Medium Intensity Statins as a Percentage of Items of All Statins by Year



Development of more proactive, preventative care models for LTC : 8 Care Processes & 3 treatment targets (all diabetes type 1 & 2)



- The percentage of people living with diabetes who have received the 8 care processes in Epping Forest North PCN is similar to the Place, but lower than ICB. However, for the three treatment targets the PCN data shows a higher percentage than Place and the ICB.
- The latest information for diabetes indicators can be found within [Ardens Manager](#).



Reduction in admissions rates of Chronic Ambulatory Care Sensitive (ACS) conditions

LTC Outcome – Reduce the rate of ambulatory care sensitive emergency hospital admissions

ICB overarching outcome of Reduce the rate of emergency admission for chronic ambulatory care sensitive conditions associated with LTCs



- Ambulatory care sensitive (ACS) conditions are conditions where effective community care and person-centred care can help prevent the need for hospital admission. (Nuffield Trust)
- Epping Forest North PCN’s admission rate for Chronic ACS conditions is lower than the ICB and Place rate, when looking at the 12 months data up to November 2023.
- AF, COPD, Heart Failure, and Diseases of the Blood, are conditions with the highest volume.
- The data on page 11 looking at observed versus expected prevalence shows an opportunity for further identification of those with COPD and Heart Failure.

Chronic Ambulatory Care Sensitive Conditions for Epping Forest North PCN	PCN Per 100,000 Rate Apr-23 to Nov-23	5 Year Trend	2024/25 Trajectory
Angina: Angina pectoris	62.39		UP
Asthma	74.79		UP
Atrial fibrillation and flutter	188.89		UP
COPD	171.99		UP
Congestive heart failure	153.79		UP
Diseases of the blood	198.11		UP
Epilepsy	44.50		UP
Hypertension	33.61		UP
Mental and behavioural disorders	11.13		UP
Nutritional, endocrine and metabolic	63.92		UP



Source: SUS Link: [Chronic ACS Conditions](#) & [NHSOF](#)

Emergency Admission rates for Falls in persons aged +75

Frailty and EoL Outcome –
Decrease rates of +75s
emergency admissions for falls
within the community

ICB overarching outcome of
Reduce the overall spend on
emergency hospital admissions

- When looking at the rolling 12 months up to November 2023 the data shows that Epping Forest North PCN has a higher rate of admissions for falls than Place, but lower rate than the ICB.
- There is variation in the data for the practices within the PCN.
- Data in the following pages shows the data for the PCN compared with Place and PCN for areas within the ECF that aim to support reducing falls. Frailty Clinical Leads will be able to advise on current programmes of work within your area aimed at reducing falls.





Emergency admission rates for Hip fractures in all over 75's

Frailty and EoL Outcome –
Decrease rates of +75s
emergency admissions for falls
within the community

ICB overarching outcome of
Reduce the overall spend on
emergency hospital admissions

- The 12 months data up to November 2023 the data shows that Epping Forest North PCN has a higher rate of admissions for hip fractures than Place and ICB.
- When looking at the data by PCN the small numbers will cause fluctuations over the years. The data for the PCN is consistently above place and ICB.
- The ECF indicators on the next page shows potential areas of opportunity through benchmarking the PCN against Place and ICB.



ECF indicators for frailty and EOL

- The data shows that Epping Forest North PCN has a lower percentage of falls frat scores completed, when compared to place and ICB as at end Dec 23.
- The percentage of the population recorded as moderately or severely frail is similar to place and higher than ICB.
- The data contained within the table below is up to the end of December, the latest position can be found at [Ardens Manager](#).

Epping Forest North Enhanced Commissioning Framework (ECF) 2023-24 for the period 1 April 23 to 31 Dec 23

	Frailty			EOL						
	Mod/Sev + falls Frat score done %	Mod frailty + SMR or polypharmacy medication review %	% Mod/Sev frailty of population	% population on EOL register	GSF %	DNACPR %	ACP %	PPD %	PPC %	SCR Consent %
ICB	16.0%	19.5%	1.9%	0.7%	64.9%	53.3%	11.3%	37.3%	39.8%	34.8%
WE	9.7%	29.0%	2.1%	0.7%	69.1%	57.1%	5.0%	33.7%	39.4%	57.6%
PCN	7.2%	44.4%	2.2%	0.9%	74.3%	46.8%	3.9%	23.0%	23.5%	61.9%
Abridge Surgery	0.0%	0.0%	1.6%	0.4%	37.5%	37.5%	0.0%	12.5%	6.3%	6.3%
High Street Surgery, Epping	0.0%	0.0%	1.0%	0.2%	84.6%	46.2%	0.0%	38.5%	46.2%	30.8%
Market Square Surgery	10.4%	51.2%	2.8%	1.2%	73.4%	45.1%	1.2%	12.7%	14.5%	80.3%
Maynard Court Surgery	5.0%	0.0%	1.3%	1.4%	63.6%	30.3%	0.8%	18.2%	16.7%	55.3%
Ongar Health Centre	13.5%	37.1%	3.4%	0.5%	92.3%	80.0%	12.3%	43.1%	41.5%	63.1%
The Limes Medical Centre	0.0%	0.0%	2.0%	1.0%	78.8%	49.5%	6.5%	28.8%	30.4%	56.0%




Hertfordshire and West Essex Integrated Care System



Prevalence of mental health conditions (QOF)


ICB overarching outcome of Improving Healthy life expectancy

- The data on this page looks at the recorded prevalence of conditions within the Mental Health section of QOF. It shows the PCN prevalence against the Place and ICB. Future iterations will include comparisons against modelled expected prevalence.
- The data shows that Epping Forest North PCN has a lower recorded prevalence for LD compared with ICB and Place which may suggest an opportunity for further identification.
- The table below shows the trend over the last three years for each area.
- The following page looks at some of the wider QOF indicators around Mental Health.



Epping Forest North PCN- Mental Health Conditions 2022-2023 QOF prevalence, with 3 year trend

	Abridge Surgery		High Street Surgery		Market Square Surgery		Maynard Court Surgery		Ongar Health Centre		The Limes Medical Centre	
	QOF Prevalence 22-23	3 year Trend	QOF Prevalence 22-23	3 year Trend	QOF Prevalence 22-23	3 year Trend	QOF Prevalence 22-23	3 year Trend	QOF Prevalence 22-23	3 year Trend	QOF Prevalence 22-23	3 year Trend
Dementia	0.9%	↘	0.6%	↘	1.2%	↗	1.7%	↘	1.0%	↘	1.4%	↗
Depression	10.7%	↗	5.9%	↗	10.6%	↘	11.3%	↗	9.9%	↘	10.6%	↗
Learning Disability	0.3%	↘	0.2%	↘	0.3%	↘	0.4%	↘	0.7%	↘	0.4%	↘
SMI	0.8%	↘	0.7%	↘	1.1%	↘	0.9%	↘	0.7%	↘	0.8%	↘



Mental Health QOF Indicators 22-23

- Mental Health QOF metrics for 2022-23 show that Epping Forest North PCN is achieving higher percentages for all metrics for both SMI and Depression in comparison to Place and the ICB.
- The data contained within the table below is the latest QOF data, the latest in year position can be found at [Ardens Manager](#).

	SMI					Depression
	% of patients with SMI who have a care plan	% of patients with SMI who have a record of BMI in the preceding 12 months	% of patients with SMI who have a record of alcohol consumption in preceding 12 months	% of patients with SMI who have a record of a lipid profile in the preceding 12 months	% of patients with SMI who have a record of blood glucose of HbA1C in preceding 12 months	% of patients with a diagnosis of depression who have been reviewed within 10-56 days
ICB	82.6	88.7	89.3	83.1	83.0	83.0
WE	77.8	86.6	87.6	82.9	82.9	79.6
Epping Forest North PCN	85.1	93.4	93.3	87.5	86.0	88.1
Abridge Surgery	92.6	96.6	100.0	92.3	91.3	100.0
High Street Surgery, Epping	80.6	97.0	100.0	93.5	96.2	85.7
Market Square Surgery	91.5	93.0	93.9	91.8	90.7	93.3
Maynard Court Surgery	94.6	98.2	94.6	85.7	82.2	87.3
Ongar Health Centre	92.4	94.2	97.1	86.8	85.9	90.1
The Limes Medical Centre	67.6	88.9	85.5	81.1	78.0	80.2





Hertfordshire and West Essex Integrated Care System



Emergency Admissions Rates for Self – Harm

ICB overarching outcome of Improving Healthy life expectancy

- Epping Forest North PCN has a lower rate of admissions for self-harm compared with both Place and ICB.
- When looking at the data it should be noted that the numbers at PCN level are small and therefore more fluctuation between the years will be seen.
- The data will continue to be monitored at wider HCP and ICB footprints.



Source: SUS